arXiv:1606.08434v1 [math.CO] 27 Jun 2016

# SUMMATION FORMULAS INVOLVING GENERALIZED HARMONIC NUMBERS

[A,B]CHUANAN WEI*, [C]XIAOXIA WANG

[A]*Department of Mathematics*
*Shanghai Normal University, Shanghai 200234, China*
[B]*Department of Information Technology*
*Hainan Medical College, Haikou 571199, China*
[C]*Department of Mathematics*
*Shanghai University, Shanghai 200444, China*

Abstract. In terms of the derivative operator and three hypergeometric series identities, several interesting summation formulas involving generalized harmonic numbers are established.

## 1. Introduction

For a complex number $x$, define the shifted-factorial to be

$$(x)_0 = 0 \quad \text{and} \quad (x)_n = x(x+1)\cdots(x+n-1) \quad \text{when} \quad n \in \mathbb{N}.$$

Following Bailey [3], define the hypergeometric series by

$${}_{1+r}F_s\left[\begin{matrix} a_0, & a_1, & \cdots, & a_r \\ & b_1, & \cdots, & b_s \end{matrix}\,\Big|\, z\right] = \sum_{k=0}^{\infty} \frac{(a_0)_k (a_1)_k \cdots (a_r)_k}{(1)_k (b_1)_k \cdots (b_s)_k} z^k,$$

where $\{a_i\}_{i\geq 0}$ and $\{b_j\}_{j\geq 1}$ are complex parameters such that no zero factors appear in the denominators of the summand on the right hand side. Then Dougall's ${}_5F_4$-series identity (cf. [2, p. 71]) can be stated as

$$\begin{aligned} &{}_5F_4\left[\begin{matrix} a, 1+\frac{a}{2}, b, c, d \\ \frac{a}{2}, 1+a-b, 1+a-c, 1+a-d \end{matrix}\,\Big|\, 1\right] \\ &= \frac{\Gamma(1+a-b)\Gamma(1+a-c)\Gamma(1+a-d)\Gamma(1+a-b-c-d)}{\Gamma(1+a)\Gamma(1+a-b-c)\Gamma(1+a-b-d)\Gamma(1+a-c-d)}, \end{aligned} \tag{1}$$

where the parameters satisfy $Re(1+a-b-c-d) > 0$ and $\Gamma(x)$ is the well-known gamma function

$$\Gamma(x) = \int_0^{\infty} t^{x-1} e^{-t} dt \ \text{ with } \ Re(x) > 0.$$

When $d = a/2$, it reduces to Dixon's ${}_3F_2$-series identity(cf. [2, p. 72]):

$$\begin{aligned} &{}_3F_2\left[\begin{matrix} a, b, c \\ 1+a-b, 1+a-c \end{matrix}\,\Big|\, 1\right] \\ &= \frac{\Gamma(1+\frac{a}{2})\Gamma(1+a-b)\Gamma(1+a-c)\Gamma(1+\frac{a}{2}-b-c)}{\Gamma(1+a)\Gamma(1+\frac{a}{2}-b)\Gamma(1+\frac{a}{2}-c)\Gamma(1+a-b-c)} \end{aligned} \tag{2}$$

*2010 Mathematics Subject Classification*: Primary 05A10 and Secondary 33C20.
*Key words and phrases.* Hypergeometric series; Derivative operator; Harmonic numbers.
*The Correspondence author*. Email addresses*: weichuanan78@163.com (C. Wei), xwang913@126.com (X. Wang).

provided that $Re(1+\frac{a}{2}-b-c)>0$. A Dixon-like identity that will appear in Section 3 is

$$\begin{aligned}
&{}_3F_2\left[\begin{matrix}a,b,c\\1+a-b,a-c\end{matrix}\,\Big|\,1\right]\\
&=\frac{1}{2^{1+c}}\frac{\Gamma(1+a-b)\Gamma(\frac{1+a}{2}-b-c)\Gamma(\frac{a-c}{2})\Gamma(\frac{1+a-c}{2})}{\Gamma(1+a-b-c)\Gamma(\frac{a}{2})\Gamma(\frac{1+a}{2}-b)\Gamma(\frac{1+a}{2}-c)}\\
&+\frac{1}{2^{1+c}}\frac{\Gamma(1+a-b)\Gamma(\frac{2+a}{2}-b-c)\Gamma(\frac{a-c}{2})\Gamma(\frac{1+a-c}{2})}{\Gamma(1+a-b-c)\Gamma(\frac{1+a}{2})\Gamma(\frac{2+a}{2}-b)\Gamma(\frac{a}{2}-c)}
\end{aligned}\tag{3}$$

provided that $Re(\frac{1+a}{2}-b-c)>0$.

For a complex number $x$ and a positive integer $\ell$, define generalized harmonic numbers of $\ell$-order to be

$$H_0^{\langle\ell\rangle}(x)=0\quad\text{and}\quad H_n^{\langle\ell\rangle}(x)=\sum_{k=1}^{n}\frac{1}{(x+k)^{\ell}}\quad\text{with}\quad n\in\mathbb{N}.$$

When $x=0$, they become harmonic numbers of $\ell$-order

$$H_0^{\langle\ell\rangle}=0\quad\text{and}\quad H_n^{\langle\ell\rangle}=\sum_{k=1}^{n}\frac{1}{k^{\ell}}\quad\text{with}\quad n\in\mathbb{N}.$$

Setting $\ell=1$ in $H_0^{\langle\ell\rangle}(x)$ and $H_n^{\langle\ell\rangle}(x)$, we obtain generalized harmonic numbers

$$H_0(x)=0\quad\text{and}\quad H_n(x)=\sum_{k=1}^{n}\frac{1}{x+k}\quad\text{with}\quad n\in\mathbb{N}.$$

When $x=0$, they reduce to classical harmonic numbers

$$H_0=0\quad\text{and}\quad H_n=\sum_{k=1}^{n}\frac{1}{k}\quad\text{with}\quad n\in\mathbb{N}.$$

For a differentiable function $f(x)$, define the derivative operator $\mathcal{D}_x$ by

$$\mathcal{D}_x f(x)=\frac{d}{dx}f(x).$$

Then it is not difficult to show that

$$\mathcal{D}_x\binom{x+s}{t}=\binom{x+s}{t}\big\{H_s(x)-H_{s-t}(x)\big\},$$

$$\mathcal{D}_x\,H_n^{\langle\ell\rangle}(x)=-\ell H_n^{\langle\ell+1\rangle}(x),$$

where $s,t\in\mathbb{N}_0$ with $t\le s$.

As pointed out by Richard Askey (cf. [1]), expressing harmonic numbers in accordance with differentiation of binomial coefficients can be traced back to Issac Newton. In 2003, Paule and Schneider [10] computed the family of series:

$$W_n(\alpha)=\sum_{k=0}^{n}\binom{n}{k}^{\alpha}\{1+\alpha(n-2k)H_k\}$$

with $\alpha=1,2,3,4,5$ by combining this way with Zeilberger's algorithm for definite hypergeometric sums. According to the derivative operator and the hypergeometric form of Andrews' $q$-series transformation, Krattenthaler and Rivoal [7] deduced general Paule-Schneider type identities with $\alpha$ being a positive integer. More results from differentiation of binomial coefficients can be found in the papers [12, 16, 17, 18]. For different ways and related harmonic number identities, the reader may refer to [4, 5, 6, 8, 9, 11, 13, 15]. It should be mentioned that Sun [14] showed recently some congruence relations concerning harmonic numbers to us.

Inspired by the work just mentioned, we shall establish several interesting summation formulas involving generalized harmonic numbers by means of the derivative operator and (1)-(3). By specifying the parameters, they can give numerous harmonic number identities. For making the reader have a taste, we select, above all, the following two ones:

$$\sum_{k=0}^{2n}\frac{(-1)^k}{\binom{2n}{k}}H_k^{\langle 2\rangle}=\frac{1+2n}{2+2n}H_n^{\langle 2\rangle},$$
$$\sum_{k=0}^{2n}\frac{(-1)^k}{\binom{2n}{k}}H_k^{2}=\frac{1+2n}{2+2n}\bigg\{H_{1+2n}^2-\frac{H_{1+2n}}{1+n}-H_{1+2n}^{\langle 2\rangle}+H_{1+n}^{\langle 2\rangle}\bigg\},$$

where the first equation comes from the case $p=0$ of (7) and the second equation is exactly Proposition 3.

## 2. Dixon's identity, reversal techniques and summation formulas involving generalized harmonic numbers

**Theorem 1.** *Let $x$ and $y$ be complex numbers. Then*

$$\begin{aligned}&\sum_{k=0}^{2n}(-1)^k\binom{2n}{k}\frac{\binom{x+k}{k}\binom{y+k}{k}}{\binom{x+2n}{k}\binom{y+2n}{k}}H_k(x)\\&=\frac{1}{2}\frac{\binom{x+n}{n}\binom{y+n}{n}\binom{1+x+y+2n}{2n}}{\binom{x+2n}{2n}\binom{y+2n}{2n}\binom{1+x+y+n}{n}}\big\{H_n(x)-H_n(1+x+y)+H_{2n}(1+x+y)\big\}.\end{aligned}$$

*Proof.* Perform the replacements $a\to -2n$, $b\to 1+x$, $c\to 1+y$ in (2) to get

$$\sum_{k=0}^{2n}(-1)^k\binom{2n}{k}\frac{\binom{x+k}{k}\binom{y+k}{k}}{\binom{x+2n}{k}\binom{y+2n}{k}}=\frac{\binom{x+n}{n}\binom{y+n}{n}\binom{1+x+y+2n}{2n}}{\binom{x+2n}{2n}\binom{y+2n}{2n}\binom{1+x+y+n}{n}}.\tag{4}$$

Applying the derivative operator $\mathcal{D}_x$ to both sides of (4), we gain

$$\begin{aligned}&\sum_{k=0}^{2n}(-1)^k\binom{2n}{k}\frac{\binom{x+k}{k}\binom{y+k}{k}}{\binom{x+2n}{k}\binom{y+2n}{k}}\{H_k(x)+H_{2n-k}(x)\}\\&=\frac{\binom{x+n}{n}\binom{y+n}{n}\binom{1+x+y+2n}{2n}}{\binom{x+2n}{2n}\binom{y+2n}{2n}\binom{1+x+y+n}{n}}\big\{H_n(x)-H_n(1+x+y)+H_{2n}(1+x+y)\big\}.\end{aligned}$$

In terms of the reversal techniques, it is not difficult to show that

$$\sum_{k=0}^{2n}(-1)^k\binom{2n}{k}\frac{\binom{x+k}{k}\binom{y+k}{k}}{\binom{x+2n}{k}\binom{y+2n}{k}}H_k(x)=\sum_{k=0}^{2n}(-1)^k\binom{2n}{k}\frac{\binom{x+k}{k}\binom{y+k}{k}}{\binom{x+2n}{k}\binom{y+2n}{k}}H_{2n-k}(x).$$

Therefore, we derive Theorem 1 to complete the proof. □

Taking $x=p$, $y=q$ with $p,q\in\mathbb{N}_0$ in Theorem 1 and utilizing (4), we achieve the summation formula on harmonic numbers:

$$\begin{aligned}&\sum_{k=0}^{2n}(-1)^k\binom{2n}{k}\frac{\binom{p+k}{k}\binom{q+k}{k}}{\binom{p+2n}{k}\binom{q+2n}{k}}H_{p+k}\\&=\frac{1}{2}\frac{\binom{p+n}{n}\binom{q+n}{n}\binom{1+p+q+2n}{2n}}{\binom{p+2n}{2n}\binom{q+2n}{2n}\binom{1+p+q+n}{n}}\big\{H_p+H_{p+n}-H_{1+p+q+n}+H_{1+p+q+2n}\big\}.\end{aligned}\tag{5}$$

**Theorem 2.** *Let $x$ be a complex number. Then*

$$\sum_{k=0}^{2n}(-1)^k\binom{2n}{k}\frac{\binom{x+k}{k}^2}{\binom{x+2n}{k}^2}H_k^{\langle 2\rangle}(x)=\frac{1}{2}\frac{\binom{x+n}{n}^2\binom{1+2x+2n}{2n}}{\binom{x+2n}{2n}^2\binom{1+2x+n}{n}}H_n^{\langle 2\rangle}(x).$$

*Proof.* Applying the derivative operator $\mathcal{D}_y$ to Theorem 1 and then fixing $y=x$, we attain

$$\begin{aligned}\sum_{k=0}^{2n}(-1)^k\binom{2n}{k}\frac{\binom{x+k}{k}^2}{\binom{x+2n}{k}^2}\{H_k^2(x)+H_k(x)H_{2n-k}(x)\}&=\frac{1}{2}\frac{\binom{x+n}{n}^2\binom{1+2x+2n}{2n}}{\binom{x+2n}{2n}^2\binom{1+2x+n}{n}}\\ \times\big\{H_n^{\langle 2\rangle}(1+2x)-H_{2n}^{\langle 2\rangle}(1+2x)+[H_n(1+2x)-H_{2n}(1+2x)-H_n(x)]^2\big\}.&\qquad(6)\end{aligned}$$

Applying the derivative operator $\mathcal{D}_x$ to Theorem 1 and then setting $y=x$, we obtain

$$\begin{aligned}&\sum_{k=0}^{2n}(-1)^k\binom{2n}{k}\frac{\binom{x+k}{k}^2}{\binom{x+2n}{k}^2}\{H_k^2(x)+H_k(x)H_{2n-k}(x)-H_k^{\langle 2\rangle}(x)\}\\ &=\frac{1}{2}\frac{\binom{x+n}{n}^2\binom{1+2x+2n}{2n}}{\binom{x+2n}{2n}^2\binom{1+2x+n}{n}}\big\{H_n^{\langle 2\rangle}(1+2x)-H_{2n}^{\langle 2\rangle}(1+2x)-H_n^{\langle 2\rangle}(x)\\ &\qquad\qquad+[H_n(1+2x)-H_{2n}(1+2x)-H_n(x)]^2\big\}.\end{aligned}$$

The difference of (6) and the last equation produces Theorem 2. □

Taking $x=p$ with $p\in\mathbb{N}_0$ in Theorem 2 and using (4), we get the summation formula on harmonic numbers:

$$\sum_{k=0}^{2n}(-1)^k\binom{2n}{k}\frac{\binom{p+k}{k}^2}{\binom{p+2n}{k}^2}H_{p+k}^{\langle 2\rangle}=\frac{1}{2}\frac{\binom{p+n}{n}^2\binom{1+2p+2n}{2n}}{\binom{p+2n}{2n}^2\binom{1+2p+n}{n}}\big\{H_{p+n}^{\langle 2\rangle}+H_p^{\langle 2\rangle}\big\}.\qquad(7)$$

**Proposition 3** (Harmonic number identity)**.**

$$\sum_{k=0}^{2n}\frac{(-1)^k}{\binom{2n}{k}}H_k^2=\frac{1+2n}{2+2n}\bigg\{H_{1+2n}^2-\frac{H_{1+2n}}{1+n}-H_{1+2n}^{\langle 2\rangle}+H_{1+n}^{\langle 2\rangle}\bigg\}.$$

*Proof.* The case $x=0$ of (6) reads as

$$\sum_{k=0}^{2n}\frac{(-1)^k}{\binom{2n}{k}}\big\{H_k^2+H_kH_{2n-k}\big\}=\frac{1+2n}{2+2n}\bigg\{\bigg(H_{1+2n}-\frac{1}{1+n}\bigg)^2-H_{1+2n}^{\langle 2\rangle}+H_{1+n}^{\langle 2\rangle}\bigg\}.$$

The case $p=n$ of Wei, Gong and Yan [17, Corollary 21] is

$$\sum_{k=0}^{2n}\frac{(-1)^k}{\binom{2n}{k}}H_kH_{2n-k}=\frac{1+2n}{2(1+n)^2}\bigg\{\frac{1}{1+n}-H_{1+2n}\bigg\}.$$

The difference of the last two equations offers Proposition 3. □

## 3. Dixon-like identity, bisection method and summation formulas

INVOLVING GENERALIZED HARMONIC NUMBERS

**Lemma 4** (Dixon-like identity)**.**

$$\begin{aligned}
&{}_3F_2\left[\begin{matrix}a,b,c\\1+a-b,a-c\end{matrix}\,\Big|\,1\right]\\
&=\frac{1}{2^{1+c}}\frac{\Gamma(1+a-b)\Gamma(\frac{1+a}{2}-b-c)\Gamma(\frac{a-c}{2})\Gamma(\frac{1+a-c}{2})}{\Gamma(1+a-b-c)\Gamma(\frac{a}{2})\Gamma(\frac{1+a}{2}-b)\Gamma(\frac{1+a}{2}-c)}\\
&+\frac{1}{2^{1+c}}\frac{\Gamma(1+a-b)\Gamma(\frac{2+a}{2}-b-c)\Gamma(\frac{a-c}{2})\Gamma(\frac{1+a-c}{2})}{\Gamma(1+a-b-c)\Gamma(\frac{1+a}{2})\Gamma(\frac{2+a}{2}-b)\Gamma(\frac{a}{2}-c)}
\end{aligned}$$

*provided that* $Re(\frac{1+a}{2}-b-c)>0$.

*Proof.* Recall Whipple's ${}_3F_2$-series identity (cf. [2, p. 149]):

$${}_3F_2\left[\begin{matrix}a,1-a,b\\c,1+2b-c\end{matrix}\,\Big|\,1\right]=\frac{\pi 2^{1-2b}\Gamma(c)\Gamma(1+2b-c)}{\Gamma(\frac{a+c}{2})\Gamma(\frac{1+a-c}{2}+b)\Gamma(\frac{1-a+c}{2})\Gamma(\frac{2-a-c}{2}+b)},$$

where $Re(b)>0$. Employ the substitution $a\to 1+a$ in the last equation to gain

$${}_3F_2\left[\begin{matrix}1+a,-a,b\\c,1+2b-c\end{matrix}\,\Big|\,1\right]=\frac{\pi 2^{1-2b}\Gamma(c)\Gamma(1+2b-c)}{\Gamma(\frac{1+a+c}{2})\Gamma(\frac{2+a-c}{2}+b)\Gamma(\frac{-a+c}{2})\Gamma(\frac{1-a-c}{2}+b)}.$$

The linear combination of the last two equations gives

$$\begin{aligned}
{}_3F_2\left[\begin{matrix}a,-a,b\\c,1+2b-c\end{matrix}\,\Big|\,1\right]&=\frac{1}{2}\frac{\pi 2^{1-2b}\Gamma(c)\Gamma(1+2b-c)}{\Gamma(\frac{a+c}{2})\Gamma(\frac{1+a-c}{2}+b)\Gamma(\frac{1-a+c}{2})\Gamma(\frac{2-a-c}{2}+b)}\\
&+\frac{1}{2}\frac{\pi 2^{1-2b}\Gamma(c)\Gamma(1+2b-c)}{\Gamma(\frac{1+a+c}{2})\Gamma(\frac{2+a-c}{2}+b)\Gamma(\frac{-a+c}{2})\Gamma(\frac{1-a-c}{2}+b)}.
\end{aligned}\tag{8}$$

According to Kummer's transformation formula (cf. [2, p. 142]):

$${}_3F_2\left[\begin{matrix}a,b,c\\d,e\end{matrix}\,\Big|\,1\right]=\frac{\Gamma(e)\Gamma(d+e-a-b-c)}{\Gamma(e-a)\Gamma(d+e-b-c)}{}_3F_2\left[\begin{matrix}a,d-b,d-c\\d,d+e-b-c\end{matrix}\,\Big|\,1\right],$$

we achieve

$$\begin{aligned}
{}_3F_2\left[\begin{matrix}c,a,b\\a-c,1+a-b\end{matrix}\,\Big|\,1\right]&=\frac{\Gamma(1+a-b)\Gamma(1+a-2b-2c)}{\Gamma(1+a-b-c)\Gamma(1+a-2b-c)}\\
&\times{}_3F_2\left[\begin{matrix}c,-c,a-b-c\\a-c,1+a-2b-c\end{matrix}\,\Big|\,1\right].
\end{aligned}$$

Evaluating the series on the right hand side by (8), we attain Lemma 4 to finish the proof. □

**Theorem 5.** *Let $x$ be a complex number. Then*

$$\begin{aligned}
&\sum_{k=0}^{n}(-1)^k\binom{n}{k}\frac{\binom{2x+k}{k}}{\binom{2x+n+k}{k}}H_k(x)\\
&=4^{n-1}\frac{\binom{n-\frac{1}{2}}{n}}{\binom{2x+2n}{n}}\big\{H_n(x)+H_n-2H_{2n}\big\}-\frac{4^{n-1}}{n}\frac{\binom{x+n-\frac{1}{2}}{n}}{\binom{2x+2n}{n}\binom{x+n}{n}}.
\end{aligned}$$

*Proof.* The case $c=-n$ of Lemma 4 can be written as

$$\begin{aligned}
&{}_3F_2\left[\begin{matrix}a,b,-n\\1+a-b,a+n\end{matrix}\,\Big|\,1\right]\\
&=2^{2n-1}\frac{(\frac{1+a}{2})_n(\frac{2+a}{2}-b)_n}{(a+n)_n(1+a-b)_n}+2^{2n-1}\frac{(\frac{a}{2})_n(\frac{1+a}{2}-b)_n}{(a+n)_n(1+a-b)_n}.
\end{aligned}$$

Replace respectively $a$ and $b$ by $1+x$ and $1+y$ in the last equation to obtain

$$\begin{aligned}&\sum_{k=0}^{n}(-1)^k\binom{n}{k}\frac{\binom{x+k}{k}\binom{y+k}{k}}{\binom{x+n+k}{k}\binom{x-y+k}{k}}\\&=\ 2^{2n-1}\frac{\binom{\frac{x}{2}+n}{n}\binom{\frac{x-1}{2}-y+n}{n}}{\binom{x+2n}{n}\binom{x-y+n}{n}}+2^{2n-1}\frac{\binom{\frac{x-1}{2}+n}{n}\binom{\frac{x-2}{2}-y+n}{n}}{\binom{x+2n}{n}\binom{x-y+n}{n}}.\end{aligned}\tag{9}$$

Applying the derivative operator $\mathcal{D}_y$ to both sides of (9), we get

$$\begin{aligned}&\sum_{k=0}^{n}(-1)^k\binom{n}{k}\frac{\binom{x+k}{k}\binom{y+k}{k}}{\binom{x+n+k}{k}\binom{x-y+k}{k}}\big\{H_k(y)+H_k(x-y)\big\}\\&=\ 2^{2n-1}\frac{\binom{\frac{x}{2}+n}{n}\binom{\frac{x-1}{2}-y+n}{n}}{\binom{x+2n}{n}\binom{x-y+n}{n}}\big\{H_n(x-y)-H_n(\tfrac{x-1}{2}-y)\big\}\\&+\ 2^{2n-1}\frac{\binom{\frac{x-1}{2}+n}{n}\binom{\frac{x-2}{2}-y+n}{n}}{\binom{x+2n}{n}\binom{x-y+n}{n}}\big\{H_n(x-y)-H_n(\tfrac{x-2}{2}-y)\big\}.\end{aligned}\tag{10}$$

In accordance with the relation

$$H_n(\tfrac{x-1}{2}-y)=2H_{2n}(x-2y)-H_n(\tfrac{x}{2}-y),$$

(10) can be reformulated as

$$\begin{aligned}&\sum_{k=0}^{n}(-1)^k\binom{n}{k}\frac{\binom{x+k}{k}\binom{y+k}{k}}{\binom{x+n+k}{k}\binom{x-y+k}{k}}\big\{H_k(y)+H_k(x-y)\big\}\\&=\ 2^{2n-1}\frac{\binom{\frac{x}{2}+n}{n}\binom{\frac{x-1}{2}-y+n}{n}}{\binom{x+2n}{n}\binom{x-y+n}{n}}\big\{H_n(x-y)+H_n(\tfrac{x}{2}-y)-2H_{2n}(x-2y)\big\}\\&+\ 2^{2n-1}\frac{\binom{\frac{x-1}{2}+n}{n}\binom{\frac{x-2}{2}-y+n}{n}}{\binom{x+2n}{n}\binom{x-y+n}{n}}\left\{H_n(x-y)-H_n(\tfrac{x}{2}-y)+\frac{2}{x-2y+2n}\right\}\\&-\ 2^{2n-1}\frac{\binom{\frac{x-1}{2}+n}{n}\binom{\frac{x}{2}-y+n}{n}}{\binom{x+2n}{n}\binom{x-y+n}{n}}\frac{2}{x-2y+2n}.\end{aligned}$$

Substitute respectively $x$ and $y$ by $2x$ and $x$ in the last equation to produce Theorem 5. □

Fixing $x=p$ with $p\in\mathbb{N}_0$ in Theorem 5 and exploiting (9), we gain the summation formula on harmonic numbers:

$$\begin{aligned}&\sum_{k=0}^{n}(-1)^k\binom{n}{k}\frac{\binom{2p+k}{k}}{\binom{2p+n+k}{k}}H_{p+k}\\&=\ 4^{n-1}\frac{\binom{n-\frac{1}{2}}{n}}{\binom{2p+2n}{n}}\big\{H_{p+n}+H_p+H_n-2H_{2n}\big\}-\frac{4^{n-1}}{n}\frac{\binom{p+n-\frac{1}{2}}{n}}{\binom{2p+2n}{n}\binom{p+n}{n}}.\end{aligned}\tag{11}$$

**Theorem 6.** *Let $x$ be a complex number. Then*

$$\begin{aligned}&\sum_{k=0}^{n}(-1)^k\binom{n}{k}\frac{\binom{\frac{x}{2}+k}{k}\binom{x-\frac{1}{2}+k}{k}}{\binom{\frac{x-1}{2}+k}{k}\binom{x-\frac{1}{2}+n+k}{k}}H_{2k}(x)\\&=\ 4^{n-1}\frac{\binom{\frac{x}{2}-\frac{1}{4}+n}{n}\binom{-\frac{3}{4}+n}{n}}{\binom{\frac{x-1}{2}+n}{n}\binom{x-\frac{1}{2}+2n}{n}}\big\{H_n(\tfrac{x-1}{2})-H_n(-\tfrac{3}{4})\big\}\\&+\ 4^{n-1}\frac{\binom{\frac{x}{2}-\frac{3}{4}+n}{n}\binom{-\frac{5}{4}+n}{n}}{\binom{\frac{x-1}{2}+n}{n}\binom{x-\frac{1}{2}+2n}{n}}\big\{H_n(\tfrac{x-1}{2})-H_n(-\tfrac{5}{4})\big\}.\end{aligned}$$

*Proof.* Perform the replacements $x \to x-\frac{1}{2}$, $y \to \frac{x}{2}$ in (10) to achieve

$$\begin{aligned}
&\sum_{k=0}^{n}(-1)^k\binom{n}{k}\frac{\binom{\frac{x}{2}+k}{k}\binom{x-\frac{1}{2}+k}{k}}{\binom{\frac{x-1}{2}+k}{k}\binom{x-\frac{1}{2}+n+k}{k}}\big\{H_k(\tfrac{x}{2})+H_k(\tfrac{x-1}{2})\big\}\\
&=\ 2^{2n-1}\frac{\binom{\frac{x}{2}-\frac{1}{4}+n}{n}\binom{-\frac{3}{4}+n}{n}}{\binom{\frac{x-1}{2}+n}{n}\binom{x-\frac{1}{2}+2n}{n}}\big\{H_n(\tfrac{x-1}{2})-H_n(-\tfrac{3}{4})\big\}\\
&+\ 2^{2n-1}\frac{\binom{\frac{x}{2}-\frac{3}{4}+n}{n}\binom{-\frac{5}{4}+n}{n}}{\binom{\frac{x-1}{2}+n}{n}\binom{x-\frac{1}{2}+2n}{n}}\big\{H_n(\tfrac{x-1}{2})-H_n(-\tfrac{5}{4})\big\}.
\end{aligned}$$

By means of the relation

$$H_k(\tfrac{x}{2})+H_k(\tfrac{x-1}{2})=2H_{2k}(x), \tag{12}$$

the last equation can be expressed as Theorem 6 to complete the proof. □

Setting $x=p$ with $p\in\mathbb{N}_0$ in Theorem 6 and utilizing (9), we attain the summation formula on harmonic numbers and generalized harmonic numbers:

$$\begin{aligned}
&\sum_{k=0}^{n}(-1)^k\binom{n}{k}\frac{\binom{\frac{p}{2}+k}{k}\binom{p-\frac{1}{2}+k}{k}}{\binom{\frac{p-1}{2}+k}{k}\binom{p-\frac{1}{2}+n+k}{k}}H_{p+2k}\\
&=\ 4^{n-1}\frac{\binom{\frac{p}{2}-\frac{1}{4}+n}{n}\binom{-\frac{3}{4}+n}{n}}{\binom{\frac{p-1}{2}+n}{n}\binom{p-\frac{1}{2}+2n}{n}}\big\{2H_p+H_n(\tfrac{p-1}{2})-H_n(-\tfrac{3}{4})\big\}\\
&+\ 4^{n-1}\frac{\binom{\frac{p}{2}-\frac{3}{4}+n}{n}\binom{-\frac{5}{4}+n}{n}}{\binom{\frac{p-1}{2}+n}{n}\binom{p-\frac{1}{2}+2n}{n}}\big\{2H_p+H_n(\tfrac{p-1}{2})-H_n(-\tfrac{5}{4})\big\}.
\end{aligned}\tag{13}$$

## 4. Dougall's identity, bisection method and summation formulas involving generalized harmonic numbers

**Theorem 7.** *Let $x$ and $y$ be complex numbers. Then*

$$\begin{aligned}
&\sum_{k=0}^{n}(-1)^k\binom{n}{k}\frac{\binom{2x+k}{k}\binom{2y+k}{k}}{\binom{2x+n+k}{k}\binom{2x-2y+k}{k}}\frac{1+2x+2k}{1+2x+n+k}H_k(x)\\
&=\ \frac{1}{2}\frac{\binom{2x+n}{n}\binom{x-2y-1+n}{n}}{\binom{x+n}{n}\binom{2x-2y+n}{n}}\big\{H_n(x)-H_n(x-2y-1)\big\}.
\end{aligned}$$

*Proof.* Employ the substitutions $a\to 1+x$, $b\to 1+y$, $c\to 1+z$ in (1) to obtain

$$\sum_{k=0}^{n}(-1)^k\binom{n}{k}\frac{\binom{x+k}{k}\binom{y+k}{k}\binom{z+k}{k}}{\binom{x+n+k}{k}\binom{x-y+k}{k}\binom{x-z+k}{k}}\frac{1+x+2k}{1+x+n+k}=\frac{\binom{x+n}{n}\binom{x-y-z-1+n}{n}}{\binom{x-y+n}{n}\binom{x-z+n}{n}}. \tag{14}$$

Applying the derivative operator $\mathcal{D}_z$ to both sides of (14), we get

$$\begin{aligned}
&\sum_{k=0}^{n}(-1)^k\binom{n}{k}\frac{\binom{x+k}{k}\binom{y+k}{k}\binom{z+k}{k}}{\binom{x+n+k}{k}\binom{x-y+k}{k}\binom{x-z+k}{k}}\frac{1+x+2k}{1+x+n+k}\big\{H_k(z)+H_k(x-z)\big\}\\
&=\ \frac{\binom{x+n}{n}\binom{x-y-z-1+n}{n}}{\binom{x-y+n}{n}\binom{x-z+n}{n}}\big\{H_n(x-z)-H_n(x-y-z-1)\big\}.
\end{aligned}\tag{15}$$

Replace respectively $x$, $y$ and $z$ by $2x$, $2y$ and $x$ in the last equation to offer Theorem 7. □

Taking $x=p$, $y=\frac{q}{2}$ with $p,q\in\mathbb{N}_0$ in Theorem 7 and using (14), we gain the summation formula on harmonic numbers:

$$\sum_{k=0}^{n}(-1)^k\binom{n}{k}\frac{\binom{2p+k}{k}\binom{q+k}{k}}{\binom{2p+n+k}{k}\binom{2p-q+k}{k}}\frac{1+2p+2k}{1+2p+n+k}H_{p+k}$$
$$=\frac{1}{2}\frac{\binom{2p+n}{n}\binom{p-q+n-1}{n}}{\binom{p+n}{n}\binom{2p-q+n}{n}}\big\{H_p+H_{p+n}+H_{p-q-1}-H_{p-q+n-1}\big\}. \tag{16}$$

**Theorem 8.** *Let $x$ and $y$ be complex numbers. Then*

$$\sum_{k=0}^{n}(-1)^k\binom{n}{k}\frac{\binom{\frac{x}{2}+k}{k}\binom{x-\frac{1}{2}+k}{k}\binom{y+k}{k}}{\binom{\frac{x-1}{2}+k}{k}\binom{x-\frac{1}{2}+n+k}{k}\binom{x-y-\frac{1}{2}+k}{k}}\frac{1+2x+4k}{1+2x+2n+2k}H_{2k}(x)$$
$$=\frac{1}{2}\frac{\binom{x-\frac{1}{2}+n}{n}\binom{\frac{x-3}{2}-y+n}{n}}{\binom{\frac{x-1}{2}+n}{n}\binom{x-y-\frac{1}{2}+n}{n}}\big\{H_n(\tfrac{x-1}{2})-H_n(\tfrac{x-3}{2}-y)\big\}.$$

*Proof.* Substitute respectively $x$ and $z$ by $x-\frac{1}{2}$ and $\frac{x}{2}$ in (15) to achieve

$$\sum_{k=0}^{n}(-1)^k\binom{n}{k}\frac{\binom{\frac{x}{2}+k}{k}\binom{x-\frac{1}{2}+k}{k}\binom{y+k}{k}}{\binom{\frac{x-1}{2}+k}{k}\binom{x-\frac{1}{2}+n+k}{k}\binom{x-y-\frac{1}{2}+k}{k}}\frac{1+2x+4k}{1+2x+2n+2k}\big\{H_k(\tfrac{x}{2})+H_k(\tfrac{x-1}{2})\big\}$$
$$=\frac{\binom{x-\frac{1}{2}+n}{n}\binom{\frac{x-3}{2}-y+n}{n}}{\binom{\frac{x-1}{2}+n}{n}\binom{x-y-\frac{1}{2}+n}{n}}\big\{H_n(\tfrac{x-1}{2})-H_n(\tfrac{x-3}{2}-y)\big\}.$$

In terms of (12), the last equation can be manipulated as Theorem 8 to finish the proof. □

Fixing $x=p$, $y=q$ with $p,q\in\mathbb{N}_0$ in Theorem 8 and exploiting (14), we attain the summation formula on harmonic numbers and generalized harmonic numbers:

$$\sum_{k=0}^{n}(-1)^k\binom{n}{k}\frac{\binom{\frac{p}{2}+k}{k}\binom{p-\frac{1}{2}+k}{k}\binom{q+k}{k}}{\binom{\frac{p-1}{2}+k}{k}\binom{p-\frac{1}{2}+n+k}{k}\binom{p-q-\frac{1}{2}+k}{k}}\frac{1+2p+4k}{1+2p+2n+2k}H_{p+2k}$$
$$=\frac{1}{2}\frac{\binom{p-\frac{1}{2}+n}{n}\binom{\frac{p-3}{2}-q+n}{n}}{\binom{\frac{p-1}{2}+n}{n}\binom{p-q-\frac{1}{2}+n}{n}}\big\{2H_p+H_n(\tfrac{p-1}{2})-H_n(\tfrac{p-3}{2}-q)\big\}. \tag{17}$$

**Theorem 9.** *Let $x$ be a complex number. Then*

$$\sum_{k=0}^{n}(-1)^k\binom{n}{k}\frac{\binom{2x+k}{k}}{\binom{2x+n+k}{k}}\frac{1+2x+2k}{1+2x+n+k}H_k^2(x)=\frac{1}{2n}\frac{\binom{2x+n}{n}}{\binom{x+n}{n}^2}\big\{H_{n-1}-H_n(x)\big\}.$$

*Proof.* Theorem 7 can be written as

$$\sum_{k=0}^{n}(-1)^k\binom{n}{k}\frac{\binom{2x+k}{k}\binom{2y+k}{k}}{\binom{2x+n+k}{k}\binom{2x-2y+k}{k}}\frac{1+2x+2k}{1+2x+n+k}H_k(x)$$
$$=\frac{1}{2(x-2y+n)}\frac{\binom{2x+n}{n}\binom{x-2y+n}{n}}{\binom{x+n}{n}\binom{2x-2y+n}{n}}\bigg\{(x-2y)\big[H_n(x)-H_n(x-2y)\big]-\frac{n}{x-2y+n}\bigg\}.$$

Applying the derivative operator $\mathcal{D}_y$ to both sides of the last equation, we obtain

$$\sum_{k=0}^{n}(-1)^k\binom{n}{k}\frac{\binom{2x+k}{k}\binom{2y+k}{k}}{\binom{2x+n+k}{k}\binom{2x-2y+k}{k}}\frac{1+2x+2k}{1+2x+n+k}H_k(x)\big\{H_k(2y)+H_k(2x-2y)\big\}$$
$$=\frac{1}{2(x-2y+n)}\frac{\binom{2x+n}{n}\binom{x-2y+n}{n}}{\binom{x+n}{n}\binom{2x-2y+n}{n}}\bigg\{(x-2y)A_n(x,y)-\frac{nB_n(x,y)}{x-2y+n}-\frac{2n}{(x-2y+n)^2}\bigg\},$$

where the corresponding symbols stand for

$$A_n(x,y) = \big[H_n(x) - H_n(x-2y)\big]\big[H_n(2x-2y) - H_n(x-2y)\big] - H_n^{\langle 2\rangle}(x-2y),$$
$$B_n(x,y) = H_n(x) + H_n(2x-2y) - 2H_n(x-2y).$$

Perform the replacement $y \to \frac{x}{2}$ in the last equation to give Theorem 9. □

Setting $x = p$ with $p \in \mathbb{N}_0$ in Theorem 9 and utilizing (14), (16) and the relation

$$H_k^2(p) = \{H_{p+k} - H_p\}^2 = H_{p+k}^2 - 2H_pH_{p+k} + H_p^2,$$

we get the summation formula on harmonic numbers:

$$\sum_{k=0}^{n}(-1)^k\binom{n}{k}\frac{\binom{2p+k}{k}}{\binom{2p+n+k}{k}}\frac{1+2p+2k}{1+2p+n+k}H_{p+k}^2 = \frac{1}{2n}\frac{\binom{2p+n}{n}}{\binom{p+n}{n}^2}\big\{H_{n-1} - H_p - H_{p+n}\big\}. \quad (18)$$

**Theorem 10.** *Let $x$ and $y$ be complex numbers. Then*

$$\begin{aligned}&\sum_{k=0}^{n}(-1)^k\binom{n}{k}\frac{\binom{\frac{x}{2}+k}{k}^2\binom{x-\frac{1}{2}+k}{k}}{\binom{\frac{x-1}{2}+k}{k}^2\binom{x-\frac{1}{2}+n+k}{k}}\frac{1+2x+4k}{1+2x+2n+2k}H_{2k}^2(x)\\ &= \frac{1}{4}\frac{\binom{x-\frac{1}{2}+n}{n}\binom{-\frac{3}{2}+n}{n}}{\binom{\frac{x-1}{2}+n}{n}^2}\Big\{\big[H_n(\tfrac{x-1}{2}) - H_n(-\tfrac{3}{2})\big]^2 - H_n^{\langle 2\rangle}(-\tfrac{3}{2})\Big\}.\end{aligned}$$

*Proof.* Applying the derivative operator $\mathcal{D}_y$ to both sides of (15), we gain

$$\begin{aligned}&\sum_{k=0}^{n}(-1)^k\binom{n}{k}\frac{\binom{x+k}{k}\binom{y+k}{k}\binom{z+k}{k}}{\binom{x+n+k}{k}\binom{x-y+k}{k}\binom{x-z+k}{k}}\frac{1+x+2k}{1+x+n+k}\\ &\times \big\{H_k(y) + H_k(x-y)\big\}\big\{H_k(z) + H_k(x-z)\big\}\\ &= \frac{\binom{x+n}{n}\binom{x-y-z-1+n}{n}}{\binom{x-y+n}{n}\binom{x-z+n}{n}}\big\{C_n(x,y) - H_n^{\langle 2\rangle}(x-y-z-1)\big\},\end{aligned}$$

where $C_n(x,y) = \big[H_n(x-y) - H_n(x-y-z-1)\big]\big[H_n(x-z) - H_n(x-y-z-1)\big]$. Employ the substitutions $x \to x - \frac{1}{2}$, $y \to \frac{x}{2}$, $z \to \frac{x}{2}$ in the last equation to achieve

$$\begin{aligned}&\sum_{k=0}^{n}(-1)^k\binom{n}{k}\frac{\binom{\frac{x}{2}+k}{k}^2\binom{x-\frac{1}{2}+k}{k}}{\binom{\frac{x-1}{2}+k}{k}^2\binom{x-\frac{1}{2}+n+k}{k}}\frac{1+2x+4k}{1+2x+2n+2k}\big\{H_k(\tfrac{x}{2}) + H_k(\tfrac{x-1}{2})\big\}^2\\ &= \frac{\binom{x-\frac{1}{2}+n}{n}\binom{-\frac{3}{2}+n}{n}}{\binom{\frac{x-1}{2}+n}{n}^2}\Big\{\big[H_n(\tfrac{x-1}{2}) - H_n(-\tfrac{3}{2})\big]^2 - H_n^{\langle 2\rangle}(-\tfrac{3}{2})\Big\}.\end{aligned}$$

According to (12), the last equation can be expressed as Theorem 10 to complete the proof. □

Taking $x = p$ with $p \in \mathbb{N}_0$ in Theorem 10 and using (14), (17) and the relation

$$H_{2k}^2(p) = \{H_{p+2k} - H_p\}^2 = H_{p+2k}^2 - 2H_pH_{p+2k} + H_p^2,$$

we attain the summation formula on harmonic numbers and generalized harmonic numbers:

$$\sum_{k=0}^{n}(-1)^k\binom{n}{k}\frac{\binom{\frac{p}{2}+k}{k}^2\binom{p-\frac{1}{2}+k}{k}}{\binom{\frac{p-1}{2}+k}{k}^2\binom{p-\frac{1}{2}+n+k}{k}}\frac{1+2p+4k}{1+2p+2n+2k}H_{p+2k}^2$$
$$=\frac{1}{4}\frac{\binom{p-\frac{1}{2}+n}{n}\binom{-\frac{3}{2}+n}{n}}{\binom{\frac{p-1}{2}+n}{n}^2}\Big\{\big[H_n(\tfrac{p-1}{2})-H_n(-\tfrac{3}{2})\big]^2-H_n^{\langle 2\rangle}(-\tfrac{3}{2})$$
$$+4H_p\big[H_p+H_n(\tfrac{p-1}{2})-H_n(-\tfrac{3}{2})\big]\Big\}. \tag{19}$$

With the change of the parameters $p$ and $q$, (5), (7), (11), (13) and (16)-(19) can create a lot of concrete harmonic number identities. The corresponding results will not be displayed here.

**Acknowledgments**

The work is supported by the National Natural Science Foundation of China (No. 11301120).

## References

[1] G. E. Andrews, K. Uchimura, Identities in combinatorics IV: differentiation and harmonic numbers, Utilitas Math. 28 (1985), 265–269.
[2] G.E. Andrews, R. Askey, R. Roy, Special Functions, Cambridge University Press, Cambridge, 2000.
[3] W.N. Bailey, Generalized Hypergeometric Series, Cambridge University Press, Cambridge, 1935.
[4] Y. Chen, Q. Hou, H. Jin, The Abel-Zeilberger algorithm, Electron. J. Comb. 18 (2011), #P17.
[5] F. Chyzak, An extension of Zeilberger's fast algorithm to general holonomic functions, Discrete Math. 217 (2000), 115–134.
[6] R.L. Graham, D.E. Knuth, O. Patashnik, Concrete Mathematics: a Foundation for Computer Science, Addison-Wesley Publishing Company, Amsterdam, 2nd edition, 1994.
[7] C. Krattenthaler, T. Rivoal, Hypergéométrie et fonction zêta de Riemann, Mem. Amer. Math. Soc. 186, no. 875, Providence, R. I., 2007.
[8] M.J. Kronenburg, Some generalized harmonic number identities, arXiv:1103.5430v2 [math.NT], 2012.
[9] M.J. Kronenburg, On two types of harmonic number identities, arXiv:1202.3981v2 [math.NT], 2012.
[10] P. Paule, C. Schneider, Computer proofs of a new family of harmonic number identities, Adv. Appl. Math. 31 (2003), 359–378.
[11] C. Schneider, Symbolic summation assists Combinatorics, Sém. Lothar. Combin. 56 (2006), Article B56b.
[12] A. Sofo, Sums of derivatives of binomial coefficients, Adv. Appl. Math. 42 (2009), 123–134.
[13] A. Sofo, Quadratic alternating harmonic number sums, J. Number Theory 154 (2015), 144–159.
[14] Z. Sun, Arithmetic theory of harmonic numbers, Proc. Amer. Math. Soc. 140 (2012), 415–428.
[15] W. Wang, Riordan arrays and harmonic number identities, Comput. Math. Appl. 60 (2010), 1494–1509.
[16] W. Wang, C. Jia, Harmonic number identities via the Newton-Andrews method, Ramanujan J. 35 (2014), 263–285.
[17] C. Wei, D. Gong, Q. Yan, Telescoping method, derivative operators and harmonic number identities, Integral Transforms Spec. Funct. 25 (2014), 203–214.
[18] J. Wang, C. Wei, Derivative operator and summation formulae involving generalized harmonic numbers, J. Math. Anal. Appl. 434 (2016) 315-341.